

THE IMPORTANCE OF IODINE FOR GOOD HEALTH

Iodine, taken in doses substantially greater than the recommended dietary allowance, has been shown to reverse fibrocystic disease of the breast and prevent breast cancer.

by Donald W. Miller, Jr, MD

© 2006–2007, LewRockwell.com

Professor of Surgery
University of Washington
Seattle, Washington, USA

Email: dwm@u.washington.edu

Websites:

<http://www.donaldmiller.com>

<http://www.LewRockwell.com>

There is growing evidence that Americans would have better health and a lower incidence of cancer and fibrocystic disease of the breast if they consumed more iodine. A decrease in iodine intake, coupled with an increased consumption of competing halogens, fluoride and bromide, has created an epidemic of iodine deficiency in America.

People in the USA consume an average of 240 micrograms (μg) of iodine a day. In contrast, people in Japan consume more than 12 milligrams (mg) (12,000 μg) of iodine a day, a 50-fold greater amount. They eat seaweed, which includes brown algae (kelp), red algae (nori sheets, with sushi) and green algae (chlorella). Compared to terrestrial plants, which contain only trace amounts of iodine (0.001 mg/g), these marine plants have high concentrations of this nutrient (0.5–8.0 mg/g). When studied in 1964, Japanese seaweed consumption was found to be 4.5 grams (g) a day, with a measured iodine concentration of 3.1 mg/g of seaweed (= 13.8 mg of iodine). According to public health officials, mainland Japanese now consume 14.5 g of seaweed a day (= 45 mg of iodine, if its iodine content, not measured, remains unchanged).¹ Researchers have determined that residents on the coast of Hokkaido eat a quantity of seaweed sufficient to provide a daily iodine intake of 200 mg a day. Saltwater fish and shellfish contain iodine, but one would have to eat 15–25 pounds of fish to get 12 mg of iodine.

Health comparisons between the two countries are disturbing. The incidence of breast cancer in the US is the highest in the world, and in Japan, until recently, is the lowest. Japanese women who emigrate from Japan or adopt a Western-style diet have a higher rate of breast cancer compared with those that consume seaweed. Life expectancy in the US is 77.85 years, 48th in 226 countries surveyed.² It is 81.25 years in Japan, the highest of all industrialised countries and only slightly behind the five leaders—Andorra, Macau, San Marino, Singapore and Hong Kong. The infant mortality rate in Japan is the lowest in the world, with 3.5 deaths under age one per 1,000 live births—half the infant mortality rate in the United States.

Today, one in seven American women (almost 15 per cent) will develop breast cancer during their lifetime. Thirty years ago, when iodine consumption was twice as high as it is now (480 μg a day), one in 20 women developed breast cancer. Iodine was used as a dough conditioner in making bread, and each slice of bread contained 0.14 mg of iodine. In 1980, breadmakers started to use bromide as a conditioner instead, which competes with iodine for absorption into the thyroid gland and other tissues in the body. Iodine was also more widely used in the dairy industry 30 years ago than it is now.

Now, iodised table salt is the chief source of iodine in a Western diet. But 45 per cent of American households buy salt without iodine, which grocery stores also sell. And over the last three decades, people who do use iodised table salt have decreased their consumption of it by 65 per cent. Furthermore, the much higher concentrations of chloride in salt (NaCl) inhibits absorption of its sister halogen iodine (the intestines absorb only 10 per cent of the iodine present in iodised table salt). As a result, 15 per cent of the US adult female population suffers from moderate to severe iodine deficiency, which health authorities define as a urinary iodine concentration less than 50 $\mu\text{g/L}$.³ Women with goitres (a visible, noncancerous enlargement of the thyroid gland) owing to iodine deficiency have been found to have a three times greater incidence of breast cancer. A high intake of iodine is associated with a low incidence in breast cancer, and a low intake with a high incidence of breast cancer.

Animal studies show that iodine prevents breast cancer, arguing for a causal association in

these epidemiological findings. The carcinogens nitrosomethylurea and DMBA cause breast cancer in more than 70 per cent of female rats. Those given iodine, especially in its molecular form as I₂, have a statistically significant decrease in incidence of cancer. Other evidence adding biologic plausibility to the hypothesis that iodine prevents breast cancer includes the finding that the ductal cells in the breast, the ones most likely to become cancerous, are equipped with an iodine pump (the sodium iodine symporter, the same one that the thyroid gland has) to soak up this element.

Similar findings apply to fibrocystic disease of the breast. The incidence of fibrocystic breast disease in American women was three per cent in the 1920s. Today, 90 per cent of women have this disorder, manifested by epithelial hyperplasia, apocrine gland metaplasia, fluid-filled cysts and fibrosis. Six million American women with fibrocystic disease have moderate to severe breast pain and tenderness that lasts more than six days during the menstrual cycle. In animal studies, female rats fed an iodine-free diet develop fibrocystic changes in their breasts, and iodine in its elemental form (I₂) cures it.

Russian researchers first showed, in 1966, that iodine effectively relieves signs and symptoms of fibrocystic breast disease. Vishniakova and Murav'eva treated 167 women suffering from fibrocystic disease with 50 mg KI (potassium iodide) during the intermenstrual period and obtained a beneficial healing effect in 71 per cent.⁴

Then Ghent and co-workers, in a study published in the *Canadian Journal of Surgery* in 1993, likewise found that iodine relieves signs and symptoms of fibrocystic breast disease in 70 per cent of their patients.⁵ This report is a composite of three clinical studies—two case series done in Canada in 696 women treated with various types of iodine, and one in Seattle. The Seattle study, done at the Virginia Mason Clinic, was a randomised, double-blind, placebo-controlled trial of 56 women, designed

to compare 3–5 mg of elemental iodine (I₂) to a placebo (an aqueous mixture of brown vegetable dye with quinine). Investigators followed the women for six months and tracked subjective and objective changes in their fibrocystic disease.

A statistical analysis of the Seattle study (enlarged to include 92 women) was done, which showed that iodine has a highly statistically significant beneficial effect on fibrocystic disease ($P < 0.001$).⁶ Iodine was shown to reduce breast tenderness, nodularity, fibrosis, turgidity and the number of macrocysts—the five parameters in a total breast examination score that a physician blinded to what treatment the woman was taking, iodine or placebo, measured. This 51-page report,⁷ now available online, was submitted to the Food and Drug Administration (FDA) in 1995, the authors seeking its approval to carry out a larger randomised controlled clinical trial on iodine for treating fibrocystic breast disease. The FDA declined to approve the study, telling lead investigator Dr Donald Low that "iodine is a natural substance, not a drug".

But the FDA has now decided to approve a similar trial, sponsored by Sympollon Pharmaceuticals. This company is

enrolling 175 women in a phase III trial, registered on ClinicalTrials.gov and sponsored by Jack Kessler, PhD.⁸

Most physicians and surgeons view iodine from a narrow perspective. It is an antiseptic that disinfects drinking water and prevents surgical wound infections, and the thyroid gland needs it to make thyroid hormones—and that's it. (When painted on the skin prior to surgery, tincture of iodine kills 90 per cent of bacteria present within 90 seconds.) The thyroid gland needs iodine to synthesise thyroxine (T₄) and triiodothyronine (T₃), hormones that regulate metabolism and steer growth and development. T₄ contains four iodine atoms combined with 27 other atoms of carbon, hydrogen, oxygen and nitrogen, but owing to its large size it accounts for 65 per cent of the molecule's weight. (T₃ has three iodine atoms.) The thyroid needs only a trace amount of iodine, 70 µg a day, to produce the requisite amount of T₄ and T₃. For that reason, thyroidologists say that iodine is best taken just in microgram amounts. They consider consuming more than 1–2 mg of iodine a day to be excessive and potentially harmful.

Expert opinion on iodine is now the purview of thyroidologists. Mainstream physicians and surgeons accept their thyroid-only view of iodine and either ignore or discount studies that show iodine in larger amounts provides extrathyroidal benefits, particularly for women's breasts. Thus a leading textbook on breast disease, Bland and Copeland's *The Breast: Comprehensive Management of Benign and Malignant Disorders* (2003), fails to mention iodine anywhere in its 1,766 pages.⁹

History, formulation and efficacy

Iodine has an important and little understood history. This relatively scarce element has played a pivotal role in the formation of our planet's atmosphere and in the evolution of life. For more than two billion years there was no oxygen in the atmosphere until a new kind of bacteria, cyanobacteria (blue-green algae), began producing

oxygen as a byproduct of photosynthesis. Cyanobacteria also developed an affinity for iodine. The most likely reason is that these organisms used iodine as an antioxidant to protect themselves against the free radicals (superoxide anion, hydrogen peroxide and hydroxyl radical) that oxygen breeds. Studying kelp, researchers have shown how iodine does this and have found that kelp will absorb increased amounts of iodine when placed under oxidative stress.¹⁰ Other researchers have shown that iodine increases the antioxidant status of human serum similar to that of vitamin C.¹¹

Iodine also induces apoptosis, programmed cell death.¹² This process is essential to growth and development (fingers form in the foetus by apoptosis of the tissue between them) and for destroying cells that represent a threat to the integrity of the organism, like cancer cells and cells infected with viruses. Human lung cancer-cells with genes spliced into them that enhance iodine uptake and utilisation undergo apoptosis and shrink when given iodine, both when grown *in vitro* outside the body and implanted in mice.¹³ Its anticancer function may well prove to be iodine's most important extrathyroidal benefit.

Iodine has other extrathyroidal functions that require more

A statistical analysis of the Seattle study... was done, which showed that iodine has a highly statistically significant beneficial effect on fibrocystic disease. Iodine was shown to reduce breast tenderness, nodularity, fibrosis, turgidity and the number of macrocysts...

study. It removes toxic chemicals¹⁴—fluoride, bromide, lead, aluminium, mercury—and biological toxins, suppresses autoimmunity,¹⁵ strengthens the T-cell adaptive immune system¹⁶ and protects against abnormal growth of bacteria in the stomach.¹⁷

In addition to the thyroid and mammary glands, other tissues possess an iodine pump (the sodium/iodine symporter). Stomach mucosa, the salivary glands and lactating mammary glands can concentrate iodine almost to the same degree as the thyroid gland (40-fold greater than its concentration in blood). Other tissues that have this pump include the ovaries, the thymus gland (seat of the adaptive immune system), the skin, the choroid plexus in the brain (which makes cerebrospinal fluid), and the joints, arteries and bones.

Today's medical establishment is wary of iodine (as it is of most naturally occurring, nonpatentable, nonpharmaceutical agents). Thyroidologists cite the Wolff–Chaikoff effect and warn that TSH (thyroid-stimulating hormone) blood levels can rise with an iodine intake of a milligram or more. The Wolff–Chaikoff effect, a temporary inhibition of thyroid hormone synthesis that supposedly occurs with increased iodine intake, is of no clinical significance.¹⁸ And an elevated

TSH, when it occurs, is "subclinical"—meaning that no signs or symptoms of hypothyroidism accompany its rise. Some people taking milligram doses of iodine, usually more than 50 mg a day, develop mild swelling of the thyroid gland without symptoms. The vast majority of people, 98–99 per cent, can take iodine in doses ranging from 10 to 200 mg a day without any clinically adverse effects on thyroid function.¹⁹ The prevalence of thyroid diseases in the 127 million people in Japan who consume high amounts of iodine is not much different than that in the USA.

Everyone agrees that a lack of iodine in the diet causes a spectrum of disorders that includes, in increasing order of severity, goitre and hypothyroidism, mental retardation and cretinism (severe mental retardation accompanied by physical deformities). Health authorities in the USA and Europe have agreed upon a Reference Daily Intake (RDI), formerly called the Recommended Dietary Allowance (RDA), for iodine, designed to prevent these disorders, which the World Health Organization (WHO) estimates afflicts 30 per cent of the world's population. The RDI for iodine, first proposed in 1980, is 100–150 µg/day. Organisations advocating this amount include the American Medical Association, the National Institutes of Health's National Research Council, the Institute of Medicine, the United Nations Food and Agricultural Organization, the WHO Expert Committee and the European Union International Programme on Chemical Safety. These health authorities consider an RDI of 100–150 µg/day of iodine sufficient to meet the requirements of nearly all (97–98 per cent) healthy individuals.

This consensus on iodine intake flies in the face of evidence justifying a higher amount. This evidence includes animal studies, *in vitro* studies on human cancer cell lines, clinical trials of iodine for fibrocystic breast disease and epidemiological data. An intake of 150 µg/day of iodine will prevent goitres and the other recognised iodine-deficiency disorders, but not breast disease. Prevention of breast disease requires higher doses of iodine. Indeed, a reasonable hypothesis is that, like goitres and cretinism, fibrocystic disease of the breast and breast cancer (and also uterine fibroids) are iodine deficiency disorders.

Indeed, a reasonable hypothesis is that, like goitres and cretinism, fibrocystic disease of the breast and breast cancer (and also uterine fibroids) are iodine-deficiency disorders.

What professor of English Albert Guerard (1914–2000) wrote about new truths applies especially to iodine: "When you seek a new path to truth, you must expect to find it blocked by expert opinion." The reigning truth on iodine is that the thyroid gland is the only organ in the body that requires this micronutrient, and a daily intake considerably more than what the thyroid gland needs is potentially harmful. The new truth is that the rest of the body also needs iodine—in milligram, not microgram, amounts. Tell that to a thyroidologist and the response will call to mind this admonition on new truths.

These are the four most common formulations of inorganic (nonradioactive) iodine, as iodide (I⁻) and with or without molecular iodine (I₂): potassium iodide (KI) tablets, in doses ranging from 0.23 to 130 mg; super-saturated potassium iodide (SSKI), 19–50 mg of iodide per drop; Lugol's solution, 6.3 mg of molecular iodine/iodide per drop; and Iodoral®, each tablet containing 12.5 mg of iodine/iodide. Both Lugol's solution and Iodoral are one-third molecular iodine (5%) and two-thirds potassium iodide (10%). Studies done to date indicate that the best iodine supplement is one that includes molecular iodine (I₂), which breast tissue prefers.

Iodine was used for a wide variety of ailments after its discovery in 1811 up until the mid-1900s, when thyroidologists warned that "excess" amounts of iodine might adversely affect thyroid function. It is effective in gram amounts for treating various dermatologic conditions, chronic lung disease, fungal infestations, tertiary syphilis and even arteriosclerosis.

The Nobel laureate Dr Albert Szent-Györgi (1893–1986), the physician who discovered vitamin C, wrote: "When I was a medical student, iodine in the form of KI was the universal medicine. Nobody knew what it did, but it did something and did something good. We students used to sum up the situation in this little rhyme:

*If ye don't know where, what and why,
Prescribe ye then K and I."*

The standard dose of potassium iodide given was one gram, which contains 770 mg of iodine.

Regarding KI and other iodine salts (like sodium iodide), the venerated 11th edition of the *Encyclopaedia Britannica*, published in 1911, states: "Their pharmacological action is as obscure as their effects in certain diseased conditions are consistently brilliant. Our ignorance of their mode of action is cloaked by the term deobstruent, which implies that they possess the power of driving out impurities from the blood and tissues. Most notably is this the case with the poisonous products of syphilis. In its tertiary stage—and also earlier—this disease yields in the most rapid and unmistakable fashion to iodides, so much so that the administration of these salts is at present the best means of determining whether, for instance, a cranial tumour be syphilitic or not."

This 19th and early 20th century medicine continues to be used in gram amounts in the 21st century by dermatologists. They treat inflammatory dermatoses, like nodular vasculitis and pyoderma gangrenosum, with SSKI, beginning with an iodine dose of 900 mg a day, followed by weekly increases up to six grams a day as tolerated. Fungal eruptions, like sporotrichosis, are treated initially in gram amounts with great success. These lesions can disappear within two weeks after treatment with iodine.

For many years, physicians used potassium iodide in doses starting at 1.5–3 grams and up to more than 10 grams a day, on and off, to treat bronchial asthma and chronic obstructive pulmonary disease with good results and surprisingly few side effects.

There is a case report in the medical literature of a 54-year-old man who, thinking it was iced tea, drank a "home preparation" of SSKI in water that his aunt kept in the refrigerator for her rheumatism. Over a short period of time he consumed 600 mL of this solution, which contained 15 g of iodide, an amount 100,000 times more than the RDI. He developed swelling of the face, neck and mouth, had transient cardiac arrhythmias and made an uneventful recovery.

The Iodine Project

Dr Guy Abraham, a former professor of obstetrics and gynaecology at UCLA, mounted what he calls "the Iodine Project" in 1997 after he read the Ghent paper on iodine for fibrocystic disease. He had his company, Optimox Corporation, make Iodoral, the tablet form of Lugol's solution, and he engaged two family-practice physicians, Dr Jorge Flechas (in 2000) in North Carolina and Dr David Brownstein (in 2003) in Michigan, to carry out clinical studies with it.

The project's hypothesis is that maintaining whole body sufficiency of iodine requires 12.5 mg a day, an amount similar to what the Japanese consume. The conventional view is that the body contains 25–50 mg of iodine, of which 70–80 per cent resides in the thyroid gland. Dr Abraham concluded that whole-body sufficiency exists when a person excretes 90 per cent of the iodine ingested. He devised an iodine-loading test where one takes 50 mg and measures the amount excreted in the urine over the next 24 hours. He found that the vast majority of people retain a substantial amount of the 50 mg dose. Many require 50 mg a day for several months before they will excrete 90 per cent of it. His studies²⁰ indicate that, given a sufficient amount, the body will retain much more iodine than originally thought—1,500 mg, with only three per cent of that amount held in the thyroid gland.

More than 4,000 patients in this project take iodine in daily doses ranging from 12.5 to 50 mg, and those with diabetes take up to 100 mg a day. These investigators have found that iodine does indeed reverse fibrocystic disease;²¹ their diabetic patients require less insulin; hypothyroid patients require less thyroid medication; symptoms of

fibromyalgia resolve; and patients with migraine headaches stop having them. To paraphrase Dr Szent-Györgi, these investigators aren't sure how iodine does it, but it does something good.

Thyroid function remained unchanged in 99 per cent of patients. Untoward effects of iodine²²—allergies, swelling of the salivary glands and the thyroid, and iodism—occurred rarely, in less than one per cent. Iodine removes the toxic halogens fluoride²³ and bromide from the body. Iodism, an unpleasant brassy taste, runny nose and acne-like skin lesions, is caused by the bromide that iodine extracts from the tissues. Symptoms subside on a lesser dose of iodine.

As these physicians point out, consuming iodine in milligram doses should, of course, be coupled with a complete nutritional program that includes adequate amounts of selenium, magnesium and omega-3 fatty acids. Done this way, an iodine intake 100 times the RDI is "the simplest, safest, most effective and least expensive way to help solve the health care crisis crippling our nation", as the leader of the Iodine Project, Dr Abraham, puts it.

People who take iodine in these amounts report that they have a greater sense of well-being, increased energy and

a lifting of brain fog. They feel warmer in cold environments, need somewhat less sleep, have an improved skin complexion and have more regular bowel movements. These purported health benefits need to be studied more thoroughly, as do those with regard to fibrocystic breast disease and cancer.

Meanwhile, perhaps we should emulate the Japanese and substantially increase our iodine intake; if not with seaweed, then with two drops of Lugol's solution (or one Iodoral tablet) a day.

About the Author:

Donald W. Miller, Jr, MD, is a cardiac surgeon and Professor of Surgery at the University of Washington in Seattle, USA. He is a member of Doctors for Disaster Preparedness and writes articles on a variety of subjects for LewRockwell.com. He can be contacted by email at dwm@u.washington.edu and via his website, <http://www.donaldmiller.com>.

Editor's Note:

To view the recommended reading list and additional endnotes attached to this article, go to the web page <http://www.lewrockwell.com/miller/miller20.html>.

The project's hypothesis is that maintaining whole-body sufficiency of iodine requires 12.5 mg a day, an amount similar to what the Japanese consume.

Endnotes

1. http://optimox.com/pics/Iodine/IOD-12/IOD_12.htm
2. <https://www.cia.gov/library/publications/the-world-factbook/>
3. <http://www.iccid.org/>
4. <http://www.donaldmiller.com/Iodine%20Talk.doc>
5. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8221402&query_hl=1&itool=pubmed_docsum
6. http://www.donaldmiller.com/Iodine_For_Fibrocystic_Disease_MX04.pdf
7. *ibid.*
8. <http://clinicaltrials.gov/ct/show/NCT00237523?order=1>
9. http://www.amazon.com/gp/product/072169490X/sr=1-1/qid=1155448365/ref=sr_1_1/104-8208774-0223107?ie=UTF8&s=books/lewrockwell/
10. [http://www.springerlink.com/\(bh4w5ojssdvwknvrhlg1tw2l\)/app/home/contribution.asp?referrer=parent&backto=issue,1,20;journal,98,977;linkingpublicationresults,1:100484,1](http://www.springerlink.com/(bh4w5ojssdvwknvrhlg1tw2l)/app/home/contribution.asp?referrer=parent&backto=issue,1,20;journal,98,977;linkingpublicationresults,1:100484,1)

11. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=10814974&query_hl=8&itool=pubmed_docsum
12. <http://www.jbc.org/cgi/content/abstract/281/28/19762>
13. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=12941836&query_hl=4&itool=pubmed_docsum
14. http://www.optimox.com/pics/Iodine/IOD-08/IOD_08.htm
15. <http://jcem.endojournals.org/cgi/reprint/81/10/3622>
16. <http://web.tiscali.it/iodio/>
17. <http://www.sciencedirect.com>, 31 October 2002
18. http://www.optimox.com/pics/Iodine/IOD-04/IOD_04.html
19. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=14057854&query_hl=6&itool=pubmed_DocSum
20. http://www.optimox.com/pics/Iodine/IOD-05/IOD_05.html
21. http://www.optimox.com/pics/Iodine/IOD-10/IOD_10.htm
22. http://www.optimox.com/pics/Iodine/IOD-09/IOD_09.htm
23. <http://www.lewrockwell.com/miller/miller17.html>